

"Express Mail" mailing label number EF 182 9660 33 US

Date of Deposit 1/31/2001

I hereby certify that this paper/fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to The Assistant Commissioner for Patents, Washington, D.C. 20231

Catherine U. Brown 44,565  
Attorney mailing application Reg No

  
Signature of Attorney mailing application

P&G Case 7996

## COMPOSITIONS AND METHODS FOR TREATING HAIR LOSS USING C<sub>16</sub> - C<sub>20</sub> AROMATIC TETRAHYDRO PROSTAGLANDINS

Mitchell Anthony deLong  
John McMillan McIver  
Robert Scott Youngquist

### FIELD OF THE INVENTION

This invention relates to compositions and methods for treating hair loss in mammals. More particularly, this invention relates to compositions and methods for arresting or reversing hair loss, or both, and promoting hair growth.

### BACKGROUND OF THE INVENTION

Hair loss is a common problem which is, for example, naturally occurring or chemically promoted through the use of certain therapeutic drugs designed to alleviate conditions such as cancer. Often such hair loss is accompanied by lack of hair re-growth which causes partial or full baldness.

Hair growth on the scalp does not occur continuously, but rather occurs by a cycle of activity involving alternating periods of growth and rest. This cycle is

divided into three main stages; anagen, catagen, and telogen. Anagen is the growth phase of the cycle and is characterized by penetration of the hair follicle deep into the dermis with rapid proliferation of cells which are differentiating to form hair.

The next phase is catagen, which is a transitional stage marked by the cessation of cell division, and during which the hair follicle regresses through the dermis and hair growth ceases. The next phase, telogen, is characterized as the resting stage during which the regressed follicle contains a germ with tightly packed dermal papilla cells. At telogen, the initiation of a new anagen phase is caused by rapid cell proliferation in the germ, expansion of the dermal papilla, and elaboration of basement membrane components. When hair growth ceases, most of the hair follicles reside in telogen and anagen is not engaged, thus causing the onset of full or partial baldness.


Attempts to invoke the re-growth of hair have been made by, for example, the promotion or prolongation of anagen. Currently, there are two drugs approved by the United States Food and Drug Administration for the treatment of male pattern baldness: topical minoxidil (marketed as ROGAINE® by Pharmacia & Upjohn), and oral finasteride (marketed as PROPECIA® by Merck & Co., Inc.). However, the search for efficacious hair growth inducers is ongoing due to factors including safety concerns and limited efficacy.

The thyroid hormone thyroxine ("T4") converts to thyronine ("T3") in human skin by deiodinase I, a selenoprotein. Selenium deficiency causes a decrease in T3 levels due to a decrease in deiodinase I activity; this reduction in T3 levels is strongly associated with hair loss. Consistent with this observation, hair growth is a reported side effect of administration of T4. See, e.g., Berman, "Peripheral Effects of L-Thyroxine on Hair Growth and Coloration in Cattle", Journal of Endocrinology, Vol. 20, pp. 282 - 292 (1960); and Gunaratnam, "The Effects of Thyroxine on Hair Growth in the Dog", J. Small Anim. Pract., Vol. 27, pp. 17 - 29 (1986). Furthermore, T3 and T4 have been the subject of several patent publications relating to treatment of hair loss. See, e.g., Fischer et al., DE 1,617,477, published January 8, 1970; Mortimer, GB 2,138,286, published October 24, 1984; and Lindenbaum, WO 96/25943, assigned to Life Medical Sciences, Inc., published August 29, 1996.

Unfortunately, however, administration of T3 or T4, or both, to treat hair loss is often not practicable because these thyroid hormones can induce significant

cardiotoxicity. See, e.g., Walker et al., U.S. Patent No. 5,284,971, assigned to Syntex, issued February 8, 1994 and Emmett et al., U.S. Patent No. 5,061,798, assigned to Smith Kline & French Laboratories, issued October 29, 1991.

In an alternative approach, prostaglandins have been proposed to promote hair growth because prostaglandins may have a similar benefit to thyroid hormones, i.e., increasing hair length and changing pigmentation. Naturally occurring prostaglandins (e.g.,  $\text{PGA}_2$ ,  $\text{PGB}_2$ ,  $\text{PGE}_1$ ,  $\text{PGF}_{2\alpha}$ , and  $\text{PGI}_2$ ) are C-20 unsaturated fatty acids.  $\text{PGF}_{2\alpha}$ , the naturally occurring Prostaglandin F analog in humans, is characterized by hydroxyl groups at the C9 and C11 positions on the alicyclic ring, a cis-double bond between C5 and C6, and a trans-double bond between C13 and C14.  $\text{PGF}_{2\alpha}$  has the formula:



Analogues of naturally occurring Prostaglandin F are known in the art. For example, see U.S. Patent No. 4,024,179 issued to Bindra and Johnson on May 17, 1977; German Patent No. DT-002,460,990 issued to Beck, Lerch, Seeger, and Teufel published on Jul. 1, 1976; U.S. Patent No. 4,128,720 issued to Hayashi, Kori, and Miyake on Dec. 5, 1978; U.S. Patent No. 4,011,262 issued to Hess, Johnson, Bindra, and Schaaf on Mar. 8, 1977; U.S. Patent No. 3,776,938 issued to Bergstrom and Sjovall on Dec. 4, 1973; P. W. Collins and S. W. Djuric, "Synthesis of Therapeutically Useful Prostaglandin and Prostacyclin Analogs", Chem. Rev., Vol. 93, pp. 1533-1564 (1993); G. L. Bundy and F. H. Lincoln, "Synthesis of 17-Phenyl-18,19,20-Trinorprostaglandins: I. The  $\text{PG}_1$  Series", Prostaglandin, Vol. 9 No. 1, pp. 1-4 (1975); W. Bartman, G. Beck, U. Lerch, H. Teufel, and B. Scholkens, "Luteolytic Prostaglandin: Synthesis and Biological Activity", Prostaglandin, Vol. 17 No. 2, pp. 301-311 (1979); C. Iljebri, G. Selen, B. Resul, J. Sternschantz, and U. Hacksell, "Derivatives of 17-Phenyl-18, 19,20-trinorprostaglandin  $\text{F}_{2\alpha}$ . Isopropyl Ester: Potential Antiglaucoma Agents", Journal of Medicinal Chemistry, Vol. 38, No. 2, pp. 289-304, (1995).

Prostaglandins in general have a wide range of biological activities. For example, PGE<sub>2</sub> has the following properties: a) regulator of cell proliferation, b) regulator of cytokine synthesis, c) regulator of immune responses and d) inducer of vasodilatation. Vasodilatation is thought to be one of the mechanisms of how

5 minoxidil provides a hair growth benefit. In vitro results in the literature also indicate some anti-inflammatory properties of the prostaglandins. c.f.; Tanaka, H., Br J. Pharm., 116, 2298, (1995).

However, previous attempts at using prostaglandins to promote hair growth have been unsuccessful. Different prostaglandin analogs can bind to multiple

10 receptors at various concentrations with a biphasic effect. Furthermore, administration of naturally occurring prostaglandins can cause side effects such as inflammation, surface irritation, smooth muscle contraction, pain, and bronchoconstriction. Therefore, it is an object of this invention to provide methods for using prostaglandin analogs to grow hair and to provide compositions that


15 promote hair growth in humans and lower animals. It is a further object of this invention to provide a selection of appropriate prostaglandin analogs that will promote hair growth and that do not cause significant undesirable side effects.

### SUMMARY OF THE INVENTION

20 This invention relates to compositions and methods for treating hair loss. The methods comprise administering the compositions comprising specific prostaglandin F analogs that interact strongly with hair-selective receptors, such as the FP receptor. The choice of prostaglandin F analog is important because it must selectively activate the FP receptor and not activate any other receptors that would negate the

25 effect of activating the FP receptor. The compositions comprise: component A) the prostaglandin F analog, component B) a carrier, and optionally component C) an activity enhancer.

Suitable prostaglandin F analogs ("PGF's") for this invention have the general formula:



wherein  $R^1$  is preferably  $\text{CO}_2\text{H}$  or  $\text{CO}_2\text{CH}_3$ ;  $R^2$  is preferably H;  $R^3$  and  $R^4$  are preferably H or  $\text{CH}_3$ ; X is preferably OH; Y is selected from the group consisting of a divalent hydrocarbon group, O, S,  $\text{S}(\text{O})$ ,  $\text{S}(\text{O})_2$ , and  $\text{NR}^{11}$ , wherein  $R^{11}$  is

5 preferably a hydrogen atom or a methyl group; and Z is preferably thienyl or phenyl. Other suitable PGF's are pharmaceutically acceptable salts, hydrates, and biohydrolyzable amides, esters, and imides of the general formula above. Optical isomers, diastereomers, and enantiomers of the structure described above are also suitable for this invention. At all stereocenters where stereochemistry is not defined

10 (i.e., C11, C12, C15, and C16), both epimers are envisioned with the epimer that corresponds to the naturally-occurring one being preferred.

### DETAILED DESCRIPTION OF THE INVENTION

This invention relates to compositions comprising prostaglandin F analogs

15 ("PGF's") to treat hair loss in mammals. "Treating hair loss" includes arresting hair loss or reversing hair loss, or both, and promoting hair growth.

Publications and patents are referred to throughout this disclosure. All U.S. Patents cited herein are hereby incorporated by reference.

All percentages, ratios, and proportions used herein are by weight unless

20 otherwise specified.

#### Definition and Usage of Terms

The following is a list of definitions for terms, as used herein:

"Activate" means binding and signal transduction of a receptor.

25 "Acyl group" means a monovalent group suitable for acylating a nitrogen atom to form an amide or carbamate, an alcohol to form a carbonate, or an oxygen atom to form an ester group. Preferred acyl groups include benzoyl, acetyl, tert-butyl

acetyl, para-phenyl benzoyl, and trifluoroacetyl. More preferred acyl groups include acetyl and benzoyl. The most preferred acyl group is acetyl.

“Aromatic group” means a monovalent group having a monocyclic ring structure or fused bicyclic ring structure. Monocyclic aromatic groups contain 5 to 10 carbon atoms, preferably 5 to 7 carbon atoms, and more preferably 5 to 6 carbon atoms in the ring. Bicyclic aromatic groups contain 8 to 12 carbon atoms, preferably 9 or 10 carbon atoms in the ring. Aromatic groups are unsubstituted. The most preferred aromatic group is phenyl.

“Carbocyclic group” means a monovalent saturated or unsaturated hydrocarbon ring. Carbocyclic groups are monocyclic, or are fused, spiro, or bridged bicyclic ring systems. Monocyclic carbocyclic groups contain 4 to 10 carbon atoms, preferably 4 to 7 carbon atoms, and more preferably 5 to 6 carbon atoms in the ring. Bicyclic carbocyclic groups contain 8 to 12 carbon atoms, preferably 9 to 10 carbon atoms in the ring. Carbocyclic groups are unsubstituted. Preferred carbocyclic groups include cyclopentyl, cyclohexyl, cyclohexenyl, cycloheptyl, and cyclooctyl. More preferred carbocyclic groups include cyclohexyl, cycloheptyl, and cyclooctyl. The most preferred carbocyclic group is cycloheptyl. Carbocyclic groups are not aromatic.

“Cyano group” means a group containing a nitrile functionality.

“FP agonist” means a compound that activates the FP receptor.

“FP receptor” means known human FP receptors, their splice variants, and undescribed receptors that have similar binding and activation profiles as the known human FP receptors. “FP” means the receptor is of the class which has the highest affinity for PGF<sub>2α</sub> of all the naturally occurring prostaglandins. FP refers to a known protein.

“Halogen atom” means F, Cl, Br, or I. Preferably, the halogen atom is F, Cl, or Br; more preferably Cl or F; and most preferably F.

“Halogenated heterogenous group” means a substituted heterogenous group or a substituted heterocyclic group, wherein at least one substituent is a halogen atom. Halogenated heterogenous groups can have a straight, branched, or cyclic structure. Preferred halogenated heterogenous groups have 1 to 12 carbon atoms,

more preferably 1 to 6 carbon atoms, and most preferably 1 to 3 carbon atoms.

Preferred halogen atom substituents are Cl and F.

“Halogenated hydrocarbon group” means a substituted monovalent hydrocarbon group or a substituted carbocyclic group, wherein at least one

- 5 substituent is a halogen atom. Halogenated hydrocarbon groups can have a straight, branched, or cyclic structure. Preferred halogenated hydrocarbon groups have 1 to 12 carbon atoms, more preferably 1 to 6 carbon atoms, and most preferably 1 to 3 carbon atoms. Preferred halogen atom substituents are Cl and F. The most preferred halogenated hydrocarbon group is trifluoromethyl.

- 10 “Heteroaromatic group” means an aromatic ring containing carbon and 1 to 4 heteroatoms in the ring. Heteroaromatic groups are monocyclic or fused bicyclic rings. Monocyclic heteroaromatic groups contain 5 to 10 member atoms (i.e., carbon and heteroatoms), preferably 5 to 7, and more preferably 5 to 6 in the ring. Bicyclic heteroaromatic rings contain 8 to 12 member atoms, preferably 9 or 10 in the ring.
- 15 Heteroaromatic groups are unsubstituted. Preferred heteroaromatic groups include thienyl, thiazolyl, purinyl, pyrimidyl, pyridyl, and furanyl. More preferred heteroaromatic groups include thienyl, furanyl, and pyridyl. The most preferred heteroaromatic group is thienyl.

- 20 “Heteroatom” means an atom other than carbon in the ring of a heterocyclic group or the chain of a heterogeneous group. Preferably, heteroatoms are selected from the group consisting of nitrogen, sulfur, and oxygen atoms. Groups containing more than one heteroatom may contain different heteroatoms.

- “Heterocyclic group” means a saturated or unsaturated ring structure containing carbon and 1 to 4 heteroatoms in the ring. No two heteroatoms are
- 25 adjacent in the ring. Heterocyclic groups are not aromatic. Heterocyclic groups are monocyclic, or are fused or bridged bicyclic ring systems. Monocyclic heterocyclic groups contain 4 to 10 member atoms (i.e., including both carbon atoms and at least 1 heteroatom), preferably 4 to 7, and more preferably 5 to 6 in the ring. Bicyclic heterocyclic groups contain 8 to 12 member atoms, preferably 9 or 10 in the ring.
- 30 Heterocyclic groups are unsubstituted. Preferred heterocyclic groups include piperzyl, morpholinyl, tetrahydrofuranyl, tetrahydropyranyl, and piperdyl.

“Heterogeneous group” means a saturated or unsaturated chain containing 1 to 18 member atoms (i.e., including both carbon and at least one heteroatom). No two heteroatoms are adjacent. Preferably, the chain contains 1 to 12 member atoms, more preferably 1 to 6, and most preferably 1 to 4. The chain may be straight or  
 5 branched. Preferred branched heterogeneous groups have one or two branches, preferably one branch. Preferred heterogeneous groups are saturated. Unsaturated heterogeneous groups have one or more double bonds, one or more triple bonds, or both. Preferred unsaturated heterogeneous groups have one or two double bonds or one triple bond. More preferably, the unsaturated heterogeneous group has one  
 10 double bond. Heterogeneous groups are unsubstituted.

“Monovalent hydrocarbon group” means a chain of 1 to 18 carbon atoms, preferably 1 to 12 carbon atoms. “Lower monovalent hydrocarbon group” means a monovalent hydrocarbon group having 1 to 6, preferably 1 to 4 carbon atoms. Monovalent hydrocarbon groups may have a straight chain or branched chain  
 15 structure. Preferred monovalent hydrocarbon groups have one or two branches, preferably 1 branch. Preferred monovalent hydrocarbon groups are saturated. Unsaturated monovalent hydrocarbon groups have one or more double bonds, one or more triple bonds, or combinations thereof. Preferred unsaturated monovalent hydrocarbon groups have one or two double bonds or one triple bond; more  
 20 preferred unsaturated monovalent hydrocarbon groups have one double bond.

“Pharmaceutically acceptable” means suitable for use in a human or other mammal.

“PGF” means a prostaglandin F analog.

“Prostaglandin” means a fatty acid derivative which has a variety of potent  
 25 biological activities of a hormonal or regulatory nature.

“Protecting group” is a group that replaces the active hydrogen of a hydroxyl moiety thus preventing undesired side reaction at the hydroxyl moiety. Use of protecting groups in organic synthesis is well known in the art. Examples of protecting groups are found in Chapter 2 Protecting Groups in Organic Synthesis by  
 30 Greene, T. W. and Wuts, P. G. M., 2<sup>nd</sup> ed., Wiley & Sons, Inc., 1991. Preferred protecting groups include silyl ethers, alkoxymethyl ethers, tetrahydropyranyl, tetrahydrofuranyl, esters, and substituted or unsubstituted benzyl ethers.



“Safe and effective amount” means a quantity of a prostaglandin high enough to provide a significant positive modification of the subject’s condition to be treated, but low enough to avoid serious side effects (at a reasonable benefit/risk ratio).

“Selective” means having a binding or activation preference for a specific  
 5 receptor over other receptors which can be quantitated based upon receptor binding or activation assays.

“Subject” means a living, vertebrate, hair- or fur-bearing animal such as a mammal (preferably human) in need of treatment.

“Substituted aromatic group” means an aromatic group wherein 1 to 4 of the  
 10 hydrogen atoms bonded to carbon atoms in the ring have been replaced with other substituents. Preferred substituents include: halogen atoms, cyano groups, monovalent hydrocarbon groups, substituted monovalent hydrocarbon groups, heterogeneous groups, aromatic groups, substituted aromatic groups, or any combination thereof. More preferred substituents include halogen atoms,  
 15 monovalent hydrocarbon groups, and substituted monovalent hydrocarbon groups. Preferred substituted aromatic groups include naphthyl. The substituents may be substituted at the ortho, meta, or para position on the ring, or any combination thereof. The preferred substitution pattern on the ring is ortho or meta. The most preferred substitution pattern is ortho.

“Substituted carbocyclic group” means a carbocyclic group wherein 1 to 4  
 20 hydrogen atoms bonded to carbon atoms in the ring have been replaced with other substituents. Preferred substituents include: halogen atoms, cyano groups, monovalent hydrocarbon groups, monovalent heterogeneous groups, substituted monovalent hydrocarbon groups, aromatic groups, substituted aromatic groups, or  
 25 any combination thereof. More preferred substituents include halogen atoms and substituted monovalent hydrocarbon groups. Carbocyclic group does not include aromatic rings.

“Substituted heteroaromatic group” means a heteroaromatic group wherein 1  
 30 to 4 hydrogen atoms bonded to carbon atoms in the ring have been replaced with other substituents. Preferred substituents include: halogen atoms, cyano groups, monovalent hydrocarbon groups, substituted monovalent hydrocarbon groups, heterogeneous groups, substituted heterogeneous groups, phenyl groups, phenoxy

groups, or any combination thereof. More preferred substituents include halogen atoms, halogenated hydrocarbon groups, halogenated heterogeneous groups, monovalent hydrocarbon groups, and phenyl groups.

“Substituted heterocyclic group” means a heterocyclic group wherein 1 to 4  
 5 hydrogen atoms bonded to carbon atoms in the ring have been replaced with other substituents. Preferred substituents include: halogen atoms, cyano groups, monovalent hydrocarbon groups, substituted monovalent hydrocarbon groups, heterogeneous groups, substituted heterogeneous groups, halogenated hydrocarbon groups, halogenated heterogeneous groups, phenyl groups, phenoxy groups, or any  
 10 combination thereof. More preferred substituents include halogen atoms and halogenated hydrocarbon groups. Substituted heterocyclic groups are not aromatic.


“Substituted heterogeneous group” means a heterogeneous group, wherein 1 to 4 of the hydrogen atoms bonded to carbon atoms in the chain have been replaced with other substituents. Preferred substituents include halogen atoms, hydroxy  
 15 groups, alkoxy groups (e.g., methoxy, ethoxy, propoxy, butoxy, and pentoxy), aryloxy groups (e.g., phenoxy, chlorophenoxy, tolyloxy, methoxyphenoxy, benzyloxy, alkyloxycarbonylphenoxy, and acyloxyphenoxy), acyloxy groups (e.g., propionyloxy, benzoyloxy, and acetoxy), carbamoyloxy groups, carboxy groups, mercapto groups, alkylthio groups, acylthio groups, arylthio groups (e.g., phenylthio,  
 20 chlorophenylthio, alkylphenylthio, alkoxyphenylthio, benzylthio, and alkyloxycarbonylphenylthio), aromatic groups (e.g., phenyl and tolyl), substituted aromatic groups (e.g., alkoxyphenyl, alkoxyphenyl, and halogenated phenyl), heterocyclic groups, heteroaromatic groups, and amino groups (e.g., amino, mono- and di- alkylamino having 1 to 3 carbon atoms, methylphenylamino,  
 25 methylbenzylamino, alkanylamido groups of 1 to 3 carbon atoms, carbamamido, ureido, and guanidino).

“Substituted monovalent hydrocarbon group” means a monovalent hydrocarbon group wherein 1 to 4 of the hydrogen atoms bonded to carbon atoms in the chain have been replaced with other substituents. Preferred substituents include  
 30 halogen atoms; halogenated hydrocarbon groups; halogenated heterogeneous groups; alkyl groups (e.g., methyl, ethyl, propyl, and butyl); hydroxy groups; alkoxy groups (e.g., methoxy, ethoxy, propoxy, butoxy, and pentoxy); aryloxy groups (e.g.,

phenoxy, chlorophenoxy, tolyloxy, methoxyphenoxy, benzyloxy, alkyloxycarbonylphenoxy, and acyloxyphenoxy); acyloxy groups (e.g., propionyloxy, benzoyloxy, and acetoxy); carbamoyloxy groups; carboxy groups; mercapto groups; alkylthio groups; acylthio groups; arylthio groups (e.g., phenylthio, chlorophenylthio, alkylphenylthio, alkoxyphenylthio, benzylthio, and alkyloxycarbonylphenylthio); aromatic groups (e.g., phenyl and tolyl); substituted aromatic groups (e.g., alkoxyphenyl, alkoxycarbonylphenyl, and halogenated phenyl); heterocyclic groups; heteroaryl groups; and amino groups (e.g., amino, mono- and di- alkylamino groups of 1 to 3 carbon atoms, methylphenylamino, methylbenzylamino, alkylamido groups of 1 to 3 carbon atoms, carbamamido, ureido, and guanidino).

#### Prostaglandin F Analogs Used in the Invention

This invention relates to the use of prostaglandin F analogs (PGFs) to treat hair loss. "Treating hair loss" means arresting hair loss, reversing hair loss, or both, and promoting hair growth. Suitable PGFs for use in this invention are selected from the group consisting of PGFs having the structure:



and pharmaceutically acceptable salts, hydrates, and biohydrolyzable amides, esters, and imides thereof.

$R^1$  is selected from the group consisting of  $CO_2H$ ,  $C(O)NHOH$ ,  $CO_2R^5$ ,  $CH_2OH$ ,  $S(O)_2R^5$ ,  $C(O)NHR^5$ ,  $C(O)NHS(O)_2R^5$ , and tetrazole.  $R^5$  is selected from the group consisting of monovalent hydrocarbon groups, substituted monovalent hydrocarbon groups, aromatic groups, substituted aromatic groups, carbocyclic groups, substituted carbocyclic groups, heterogeneous groups, substituted heterogeneous groups, heterocyclic groups, substituted heterocyclic groups, heteroaromatic groups, and substituted heteroaromatic groups. Preferably,  $R^5$  is selected from the group consisting of  $CH_3$ ,  $C_2H_5$ , and  $C_3H_7$ . Preferably,  $R^1$  is

selected from the group consisting of  $\text{CO}_2\text{H}$ ,  $\text{CO}_2\text{CH}_3$ ,  $\text{CO}_2\text{C}_2\text{H}_5$ ,  $\text{CO}_2\text{C}_3\text{H}_7$ ,  $\text{CO}_2\text{C}_4\text{H}_9$ ,  $\text{CO}_2\text{C}_3\text{H}_7\text{O}_2$ , and  $\text{C}(\text{O})\text{NHS}(\text{O})_2\text{R}^5$ . More preferably,  $\text{R}^1$  is selected from the group consisting of  $\text{CO}_2\text{H}$ ,  $\text{CO}_2\text{CH}_3$ ,  $\text{CO}_2\text{C}_2\text{H}_5$ , and  $\text{CO}_2\text{C}_3\text{H}_7$ . Most preferably,  $\text{R}^1$  is selected from the group consisting of  $\text{CO}_2\text{H}$ ,  $\text{CO}_2\text{CH}_3$ , and  $\text{CO}_2\text{C}_3\text{H}_7$ .

5  $\text{R}^2$  is a hydrogen atom or a lower monovalent hydrocarbon group.  $\text{R}^2$  is preferably a hydrogen atom or a methyl group.

$\text{R}^3$  and  $\text{R}^4$  are each independently selected from the group consisting of H,  $\text{CH}_3$ ,  $\text{C}_2\text{H}_5$ ,  $\text{OR}^{10}$ ,  $\text{SR}^{10}$ , and OH; with the proviso that both  $\text{R}^3$  and  $\text{R}^4$  are not OH.  $\text{R}^{10}$  is selected from the group consisting of a monovalent hydrocarbon group, a substituted monovalent hydrocarbon group, a heterogeneous group, a substituted heterogeneous group, a carbocyclic group, a substituted carbocyclic group, an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group; with the proviso that  $\text{R}^{10}$  has 1 to 8 member atoms. Preferably,  $\text{R}^3$  and  $\text{R}^4$  are both hydrogen atoms.

15 X is selected from the group consisting of  $\text{NR}^6\text{R}^7$ ,  $\text{OR}^8$ ,  $\text{SR}^9$ ,  $\text{S}(\text{O})\text{R}^9$ , and  $\text{S}(\text{O})_2\text{R}^9$ . Preferably, X is selected from the group consisting of  $\text{NR}^6\text{R}^7$  and  $\text{OR}^8$ . More preferably, X is OH.

$\text{R}^6$ ,  $\text{R}^7$ , and  $\text{R}^8$  are each independently selected from the group consisting of hydrogen atoms, acyl groups, monovalent hydrocarbon groups, substituted monovalent hydrocarbon groups, heterogeneous groups, substituted heterogeneous groups, carbocyclic groups, substituted carbocyclic groups, aromatic groups, substituted aromatic groups, heteroaromatic groups, and substituted heteroaromatic groups. Preferably,  $\text{R}^6$  and  $\text{R}^7$  are selected from the group consisting of H,  $\text{CH}_3$ , and  $\text{C}_2\text{H}_5$ . Preferably,  $\text{R}^8$  is selected from the group consisting of H,  $\text{CH}_3$ ,  $\text{C}_2\text{H}_5$ , and  $\text{C}_3\text{H}_7$ .

$\text{R}^9$  is selected from the group consisting of monovalent hydrocarbon groups, substituted monovalent hydrocarbon groups, heterogeneous groups, substituted heterogeneous groups, carbocyclic groups, substituted carbocyclic groups, heterocyclic groups, substituted heterocyclic groups, aromatic groups, substituted aromatic groups, heteroaromatic groups, and substituted heteroaromatic groups. Preferably,  $\text{R}^9$  is selected from the group consisting of  $\text{CH}_3$ , and  $\text{C}_2\text{H}_5$ .

Y is selected from the group consisting of an oxygen atom, a divalent hydrocarbon group, a sulfur-containing moiety, and a nitrogen-containing group. The divalent hydrocarbon group has the formula  $(CH_2)_n$ , wherein n is an integer with a value of 0 to 3. Preferably, n is 0, 1, or 2; more preferably, n is 1.

5           The sulfur-containing moiety for Y is selected from the group consisting of a sulfur atom, S(O), and S(O)<sub>2</sub>. When Y is a sulfur-containing moiety, it preferably is a sulfur atom.

          The nitrogen-containing group for Y has the formula NR<sup>11</sup>. R<sup>11</sup> is selected from the group consisting of a hydrogen atom, an acyl group, a monovalent  
10   hydrocarbon group, a substituted monovalent hydrocarbon group, a heterogeneous group, a substituted heterogeneous group, a carbocyclic group, a substituted carbocyclic group, a heterocyclic group, a substituted heterocyclic group, an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group. Preferably R<sup>11</sup> is H or CH<sub>3</sub>.

15           Z is selected from the group consisting of a carbocyclic group, a substituted carbocyclic group, a heterocyclic group, a substituted heterocyclic group, an aromatic group, a substituted aromatic group, a heteroaromatic group, and a substituted heteroaromatic group. Preferably Z is selected from the group consisting of a monocyclic carbocyclic group, a substituted monocyclic carbocyclic group, a  
20   monocyclic heterocyclic group, a substituted monocyclic heterocyclic group, a monocyclic aromatic group, a substituted monocyclic aromatic group, a monocyclic heteroaromatic group, and a substituted monocyclic heteroaromatic group. More preferably, Z is selected from the group consisting of a monocyclic aromatic group, a substituted monocyclic aromatic group, a monocyclic heteroaromatic group, and a  
25   substituted monocyclic heteroaromatic group. Most preferably, Z is thienyl or phenyl.

          Optical isomers, diastereomers, and enantiomers of the structure described above are also suitable for use in this invention. At all stereocenters where stereochemistry is not defined (i.e., C11, C12, C15, and C16), both epimers are  
30   envisioned.

          Examples of suitable PGF's having the formula above wherein Y is selected from the group consisting of NR<sup>11</sup>, S, S(O), and S(O)<sub>2</sub> include:

- 13,14-dihydro-16-(3-methylphenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;  
 13,14-dihydro-16-(3-methylphenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α;  
 3,14-dihydro-16-(3-fluorophenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α;  
 13,14-dihydro-16-(3-fluorophenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;  
 5 13,14-dihydro-16-(3-fluorophenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α isopropyl ester;  
 13,14-dihydro-16-(2,6-difluorophenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;  
 13,14-dihydro-16-(3,5-difluorophenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α;  
 10 13,14-dihydro-16-(2-methylphenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;  
 13,14-dihydro-16-(4-methylphenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;  
 13,14-dihydro-16-(4-methylphenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α;  
 13,14-dihydro-16-(2-fluorophenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;  
 13,14-dihydro-16-(2-fluorophenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α;  
 15 13,14-dihydro-15-methyl-16-(3-fluorophenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;  
 13,14-dihydro-15-methyl-16-(3-fluorophenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α;  
 13,14-dihydro-15-methyl-16-(2-methylphenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;  
 20 13,14-dihydro-15-methyl-16-(2-methylphenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α;  
 13,14-dihydro-16-(3-fluorophenylsulfonyl)-16-tetranor Prostaglandin F<sub>1</sub>α;  
 13,14-dihydro-16-(3-methylphenylamino)-16-tetranor prostaglandin F<sub>1</sub>α methyl ester;  
 13,14-dihydro-16-(3-methylphenylamino)-16-tetranor prostaglandin F<sub>1</sub>α;  
 25 13,14-dihydro-16-(2-methylphenylamino)-16-tetranor prostaglandin F<sub>1</sub>α methyl ester;  
 13,14-dihydro-16-(2-methylphenylamino)-16-tetranor prostaglandin F<sub>1</sub>α;  
 13,14-dihydro-16-(2-fluorophenylthio)-16-tetranor prostaglandin F<sub>1</sub>α 1-hydroxamic acid;

- 13,14-dihydro-16-(3-chlorophenylamino)-16-tetranor prostaglandin F<sub>1</sub>α 1-hydroxamic acid;
- 13,14-dihydro-16-(3-trifluoromethylphenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;
- 5 13,14-dihydro-16-(3-trifluoromethylphenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α;
- 13,14-dihydro-16-(3-trifluoromethylphenylthio)-16-tetranor prostaglandin F<sub>1</sub>α 1-hydroxamic acid;
- 13,14-dihydro-16-(phenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;
- 13,14-dihydro-16-(phenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α isopropyl ester;
- 10 13,14-dihydro-15-methyl-16-(phenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;
- 13,14-dihydro-15-methyl-16-(phenylthio)-16-tetranor Prostaglandin F<sub>1</sub>α;
- 13,14-dihydro-16-(phenylamino)-16-tetranor prostaglandin F<sub>1</sub>α methyl ester;
- 13,14-dihydro-16-(phenylamino)-16-tetranor prostaglandin F<sub>1</sub>α;
- 15 13,14-dihydro-16-(2-thienylthio)-16-tetranor prostaglandin F<sub>1</sub>α methyl ester;
- 13,14-dihydro-16-(2-thienylthio)-16-tetranor prostaglandin F<sub>1</sub>α;
- 13,14-dihydro-16-(1-naphthylthio)-16-tetranor Prostaglandin F<sub>1</sub>α isopropyl ester;
- 13,14-dihydro-16-(1-naphthylthio)-16-tetranor Prostaglandin F<sub>1</sub>α; and
- 13,14-dihydro-15-butoxy-15-dehydroxy-16-(phenylthio)-16-tetranor prostaglandin
- 20 F<sub>1</sub>α methyl ester.

Examples of suitable PGF's having the formula above wherein Y is a divalent hydrocarbon group include:

- 13,14-dihydro-17-(2,4-difluorophenyl)-17-trinor prostaglandin F<sub>1</sub>α methyl ester;
- 13,14-dihydro-17-(2,4-difluorophenyl)-17-trinor prostaglandin F<sub>1</sub>α;
- 25 13,14-dihydro-17-(2-fluorophenyl)-17-trinor prostaglandin F<sub>1</sub>α methyl ester;
- 13,14-dihydro-17-(2-fluorophenyl)-17-trinor prostaglandin F<sub>1</sub>α;
- 13,14-dihydro-17-(3-fluorophenyl)-17-trinor prostaglandin F<sub>1</sub>α methyl ester;
- 13,14-dihydro-17-(3-fluorophenyl)-17-trinor prostaglandin F<sub>1</sub>α;
- 13,14-dihydro-17-(4-fluorophenyl)-17-trinor prostaglandin F<sub>1</sub>α methyl ester;

- 13,14-dihydro-17-(4-fluorophenyl)-17-trinor prostaglandin  $F_1\alpha$ ;
- 13,14-dihydro-17-(2-methoxyphenyl)-17-trinor prostaglandin  $F_1\alpha$  methyl ester;
- 13,14-dihydro-17-(2-methoxyphenyl)-17-trinor prostaglandin  $F_1\alpha$ ;
- 13,14-dihydro-17-(3-methoxyphenyl)-17-trinor prostaglandin  $F_1\alpha$  methyl ester;
- 5 13,14-dihydro-17-(3-methoxyphenyl)-17-trinor prostaglandin  $F_1\alpha$ ;
- 13,14-dihydro-17-(4-methoxyphenyl)-17-trinor prostaglandin  $F_1\alpha$  methyl ester;
- 13,14-dihydro-17-(4-methoxyphenyl)-17-trinor prostaglandin  $F_1\alpha$ ;
- 13,14-dihydro-17-(3,5-difluorophenyl)-17-trinor prostaglandin  $F_1\alpha$  isopropyl ester;
- 13,14-dihydro-18-(2-thienyl)-18-dinor prostaglandin  $F_1\alpha$  methyl ester;
- 10 13,14-dihydro-18-(2-thienyl)-18-dinor prostaglandin  $F_1\alpha$  isopropyl ester;
- 13,14-dihydro-17-((2-trifluoromethyl)phenyl)-17-trinor prostaglandin  $F_1\alpha$  methyl ester;
- 13,14-dihydro-17-((2-trifluoromethyl)phenyl)-17-trinor prostaglandin  $F_1\alpha$ ;
- 13,14-dihydro-17-((3-trifluoromethyl)phenyl)-17-trinor prostaglandin  $F_1\alpha$  methyl ester;
- 13,14-dihydro-17-((3-trifluoromethyl)phenyl)-17-trinor prostaglandin  $F_1\alpha$ ;
- 15 13,14-dihydro-17-((4-trifluoromethyl)phenyl)-17-trinor prostaglandin  $F_1\alpha$  methyl ester;
- 13,14-dihydro-17-((4-trifluoromethyl)phenyl)-17-trinor prostaglandin  $F_1\alpha$ ;
- 13,14-dihydro-17-(2-methylphenyl)-17-trinor prostaglandin  $F_1\alpha$  methyl ester;
- 13,14-dihydro-17-(2-methylphenyl)-17-trinor prostaglandin  $F_1\alpha$ ;
- 13,14-dihydro-17-(3-methylphenyl)-17-trinor prostaglandin  $F_1\alpha$  methyl ester;
- 20 13,14-dihydro-17-(3-methylphenyl)-17-trinor prostaglandin  $F_1\alpha$ ;
- 13,14-dihydro-17-(4-methylphenyl)-17-trinor prostaglandin  $F_1\alpha$  methyl ester; and
- 13,14-dihydro-17-(4-methylphenyl)-17-trinor prostaglandin  $F_1\alpha$ .

Examples of suitable PGF's having the formula above wherein Y is an oxygen atom include:

- 25 13,14-dihydro-16,16-dimethyl-16-(2-fluorophenoxy)-16-tetranor prostaglandin  $F_1\alpha$ ;
- 13,14-dihydro-16,16-dimethyl-16-(2-methylphenoxy)-16-tetranor Prostaglandin  $F_1\alpha$ ;
- 13,14-dihydro-16,16-dimethyl-16-(2,3 difluorophenoxy)-16-tetranor Prostaglandin  $F_1\alpha$ ;
- 13,14-dihydro-16-(2,5 difluorophenoxy)-16-tetranor Prostaglandin  $F_1\alpha$ ;



- 13,14-dihydro-16-(3-fluoro-5-trifluoromethyl phenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α isopropyl ester;
- 13,14-dihydro-16,16-dimethyl-16-(4-chlorophenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α;
- 13,14-dihydro-16-methyl-16-(3-chlorophenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α isopropyl ester;
- 13,14-dihydro-16-isopropyl-16-(2-fluorophenoxy)-16-tetranor prostaglandin F<sub>1</sub>α;
- 13,14-dihydro-16-ethyl-16-(2-methylphenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α isopropyl ester;
- 13,14-dihydro-16-(hydroxymethyl)-16-phenoxy-16-tetranor Prostaglandin F<sub>1</sub>α;
- 13,14-dihydro-16-methyl-16-(4-ethylphenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α;
- 13,14-dihydro-16-methyl-16-(3-chlorophenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;
- 13,14-dihydro-16-methyl-16-(4-phenylphenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α isopropyl ester;
- 13,14-dihydro-16,16-dimethyl-16-(4-phenoxyphenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α isopropyl ester;
- 13,14-dihydro-16,16-dimethyl-16-(2-fluorophenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α hydroxamic acid;
- 13,14-dihydro-16-methyl-16-(3-chlorophenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α 1-hydroxamic acid;
- 13,14-dihydro-16-methoxymethyl-16-(2,3-difluorophenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α 1-hydroxamic acid;
- 13,14-dihydro-16-phenoxy-16-tetranor Prostaglandin F<sub>1</sub>α methanesulfonamide;
- 13,14-dihydro-15-fluoro-16-(2-fluorophenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;
- 13,14-dihydro-15-methyl-16,16-dimethyl-16-(2-fluorophenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α;
- 13, 14-dihydro-15-fluoro-16-(2,3-difluorophenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α 1-hydroxamic acid;
- 13,14-dihydro-15-methylthio-15-dehydroxy-16-(2-methylphenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α;

13,14-dihydro-15-methylthio-15-dehydroxy-16-methyl-16-(2-methylphenoxy) 16-tetranor Prostaglandin F<sub>1</sub>α 1-hydroxamic acid;

13,14-dihydro-15-methoxy-16,16-dimethyl-16-(2-fluorophenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α 1-hydroxamic acid;

5 13,14-dihydro-15-ethoxy-15-dehydroxy-16-phenoxy-16-tetranor Prostaglandin F<sub>1</sub>α isopropyl ester;

13,14-dihydro-15-sulfonylmethyl-15-dehydroxy-16-methyl-16-(2-methylphenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;

10 13,14-dihydro-15-sulfoxymethyl-15-dehydroxy-16-methyl-16-(2-methylphenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;

13,14-dihydro-15-methyl-15-methylamino-15-dehydroxy-16,16-dimethyl-16-(2-fluorophenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α methyl ester;

13,14-dihydro-15-methyl-15-methylamino-15-dehydroxy-16-methyl-16-(2-methylphenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α 1-hydroxamic acid;

15 13,14-dihydro-15-methyl-15-(N,N-dimethylamino)-16-ethyl-16-(2-fluorophenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α isopropyl ester; and

13,14-dihydro-16(2,6-difluorophenoxy)-16-tetranor Prostaglandin F<sub>1</sub>α glyceryl ester.

Suitable PGF's are known in the art. Examples of suitable PGF's and methods for their preparation are disclosed in International Published Patent Application Numbers WO  
20 99/12895A1, WO 99/12896A1, and WO 99/12899A1, and in U.S. Patent No. 5,977,173.

### Compositions of the Invention

This invention further relates to a composition for treating hair loss. The composition comprises A) the PGF described above and B) a carrier. The  
25 composition may further comprise C) one or more optional activity enhancers. The composition can be a pharmaceutical or cosmetic composition, administered for treatment or prophylaxis of hair loss. Standard pharmaceutical formulation techniques are used, such as those disclosed in Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa (1990).

The composition further comprises component B) a carrier. "Carrier" means one or more compatible substances that are suitable for administration to a mammal. Carrier includes solid or liquid diluents, hydrotopes, surface-active agents, and encapsulating substances. "Compatible" means that the components of the composition are capable of being commingled with the PGF's, and with each other, in a manner such that there is no interaction which would substantially reduce the efficacy of the composition under ordinary use situations. Carriers must be of sufficiently high purity and sufficiently low toxicity to render them suitable for administration to the mammal being treated. The carrier can be inert, or it can possess pharmaceutical benefits, cosmetic benefits, or both.

The choice of carrier for component B) depends on the route by which A) the PGF will be administered and the form of the composition. The composition may be in a variety of forms, suitable, for example, for systemic administration (e.g., oral, rectal, nasal, sublingual, buccal, or parenteral) or topical administration directly to the locus of desired hair growth (e.g., local application on the skin, ocular, liposome delivery systems, or iontophoresis). Topical administration is preferred.

Carriers for systemic administration typically comprise one or more ingredients selected from the group consisting of a) diluents, b) lubricants, c) binders, d) disintegrants, e) colorants, f) flavors, g) sweeteners, h) antioxidants, j) preservatives, k) glidants, m) solvents, n) suspending agents, o) surfactants, combinations thereof, and others.

Ingredient a) is a diluent. Suitable diluents include sugars such as glucose, lactose, dextrose, and sucrose; polyols such as propylene glycol; calcium carbonate; sodium carbonate; glycerin; mannitol; and sorbitol.

Ingredient b) is a lubricant. Suitable lubricants are exemplified by solid lubricants including silica, talc, stearic acid and its magnesium salts and calcium salts, calcium sulfate; and liquid lubricants such as polyethylene glycol and vegetable oils such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil and oil of theobroma.

Ingredient c) is a binder. Suitable binders include polyvinylpyrrolidone; magnesium aluminum silicate; starches such as corn starch and potato starch;

gelatin; tragacanth; and cellulose and its derivatives, such as ethyl cellulose, methylcellulose, hydroxypropylmethylcellulose, microcrystalline cellulose, and sodium carboxymethylcellulose; carbomer; providone; acacia; guar gum; and xanthan gum.

- 5           Ingredient d) is a disintegrant. Suitable disintegrants include agar, alginic acid and the sodium salt thereof, effervescent mixtures, croscarmellose, crospovidone, sodium carboxymethyl starch, sodium starch glycolate, clays, and ion exchange resins.

          Ingredient e) is a colorant such as an FD&C dye.

- 10          Ingredient f) is a flavor such as menthol, peppermint, and fruit flavors.

          Ingredient g) is a sweetener such as aspartame and saccharin.

          Ingredient h) is an antioxidant such as butylated hydroxyanisole, butylated hydroxytoluene, and vitamin E.

- Ingredient j) is a preservative such as phenol, alkyl esters of  
15       parahydroxybenzoic acid, benzoic acid and the salts thereof, boric acid and the salts thereof, sorbic acid and the salts thereof, chlorbutanol, benzyl alcohol, thimerosal, phenylmercuric acetate and nitrate, nitromersol, benzalkonium chloride, cetylpyridinium chloride, methyl paraben, and propyl paraben. Particularly preferred are the salts of benzoic acid, cetylpyridinium chloride, methyl paraben and propyl  
20       paraben, and sodium benzoate.

          Ingredient k) is a glidant such as silicon dioxide.

- Ingredient m) is a solvent, such as water, isotonic saline, ethyl oleate, alcohols such as ethanol, glycerin, glycols (e.g., polypropylene glycol and polyethylene glycol), and buffer solutions (e.g., phosphate, potassium acetate, boric  
25       carbonic, phosphoric, succinic, malic, tartaric, citric, acetic, benzoic, lactic, glyceric, gluconic, glutaric and glutamic).

          Ingredient n) is a suspending agent. Suitable suspending agents include AVICEL® RC-591 from FMC Corporation of Philadelphia, Pennsylvania and sodium alginate.

- 30          Ingredient o) lecithin, polysorbate 80, sodium lauryl sulfate, polyoxyethylene sorbitan fatty acid esters, polyoxyethylene monoalkyl ethers, sucrose monoesters, lanolin esters, and lanolin ethers. Suitable surfactants are known in the art and

commercially available, e.g., the TWEENS® from Atlas Powder Company of Wilmington, Delaware.

Compositions for parenteral administration typically comprise A) 0.1 to 10% of a PGF and B) 90 to 99.9% of a carrier comprising a) a diluent and m) a solvent.

- 5 Preferably, component a) is propylene glycol and m) is ethanol or ethyl oleate.

Compositions for oral administration can have various dosage forms. For example, solid forms include tablets, capsules, granules, and bulk powders. These oral dosage forms comprise a safe and effective amount, usually at least 5%, and preferably from 25% to 50%, of A) the PGF. The oral dosage

- 10 compositions further comprise B) 50 to 95% of a carrier, preferably 50 to 75%.

- Tablets can be compressed, tablet triturates, enteric-coated, sugar-coated, film-coated, or multiple-compressed. Tablets typically comprise A) the PGF, and B) a carrier comprising ingredients selected from the group consisting of a) diluents, b) lubricants, c) binders, d) disintegrants, e) colorants, f) flavors, g) sweeteners, k) glidants, and combinations thereof. Preferred diluents include calcium carbonate, sodium carbonate, mannitol, lactose and cellulose. Preferred binders include starch, gelatin, and sucrose. Preferred disintegrants include alginic acid, and croscarmellose. Preferred lubricants include magnesium stearate, stearic acid, and talc. Preferred colorants are the FD&C dyes, which can be added for appearance. Chewable tablets preferably contain g) sweeteners such as aspartame and saccharin, or f) flavors such as menthol, peppermint, and fruit flavors.
- 15  
20

- Capsules (including time release and sustained release formulations) typically comprise A) the PGF, and B) a carrier comprising one or more a) diluents disclosed above in a capsule comprising gelatin. Granules typically comprise A) the PGF, and preferably further comprise k) glidants such as silicon dioxide to improve flow characteristics.
- 25

- The selection of ingredients in the carrier for oral compositions depends on secondary considerations like taste, cost, and shelf stability, which are not critical for the purposes of this invention. One skilled in the art can optimize appropriate ingredients without undue experimentation.
- 30

The solid compositions may also be coated by conventional methods, typically with pH or time-dependent coatings, such that A) the PGF is released in the

gastrointestinal tract at various times to extend the desired action. The coatings typically comprise one or more components selected from the group consisting of cellulose acetate phthalate, polyvinylacetate phthalate, hydroxypropyl methyl cellulose phthalate, ethyl cellulose, EUDRAGIT® coatings (available from Rohm & Haas G.M.B.H. of Darmstadt, Germany), waxes and shellac.

Compositions for oral administration can also have liquid forms. For example, suitable liquid forms include aqueous solutions, emulsions, suspensions, solutions reconstituted from non-effervescent granules, suspensions reconstituted from non-effervescent granules, effervescent preparations reconstituted from effervescent granules, elixirs, tinctures, syrups, and the like. Liquid orally administered compositions typically comprise A) the PGF and B) a carrier comprising ingredients selected from the group consisting of a) diluents, e) colorants, and f) flavors, g) sweeteners, j) preservatives, m) solvents, n) suspending agents, and o) surfactants. *Peroral* liquid compositions preferably comprise one or more ingredients selected from the group consisting of e) colorants, f) flavors, and g) sweeteners.

Other compositions useful for attaining systemic delivery of the subject compounds include sublingual, buccal and nasal dosage forms. Such compositions typically comprise one or more of soluble filler substances such as a) diluents including sucrose, sorbitol and mannitol; and c) binders such as acacia, microcrystalline cellulose, carboxymethyl cellulose, and hydroxypropyl methyl cellulose. Such compositions may further comprise b) lubricants, e) colorants, f) flavors, g) sweeteners, h) antioxidants, and k) glidants.

The compositions may further comprise component C) an optional activity enhancer. Component C) is preferably selected from the group consisting of i) hair growth stimulants (other than the PGF) and ii) penetration enhancers.

Component i) is an optional hair growth stimulant. Component i) is exemplified by vasodilators, antiandrogens, cyclosporins, cyclosporin analogs, antimicrobials, anti-inflammatories, thyroid hormones, thyroid hormone derivatives, and thyroid hormone analogs, non-selective prostaglandin agonists or antagonists, retinoids, triterpenes, combinations thereof, and others. "Non-selective

prostaglandin" agonists and antagonists differ from component A) in that they do not selectively activate the FP receptor, and they may activate other receptors.

Vasodilators such as potassium channel agonists including minoxidil and minoxidil derivatives such as aminexil and those described in U.S. Patent Numbers  
 5 3,382,247, 5,756,092, 5,772,990, 5,760,043, 5,466,694, 5,438,058, 4,973,474, and cromakalin and diazoxide can be used as optional hair growth stimulants in the composition.

Examples of suitable antiandrogens include 5- $\alpha$ -reductase inhibitors such as finasteride and those described in U.S. Patent Number 5,516,779, and in Nane et al.,  
 10 Cancer Research 58, "Effects of Some Novel Inhibitors of C17,20-Lyase and 5 $\alpha$ -Reductase *in vitro* and *in vivo* and Their Potential Role in the Treatment of Prostate Cancer," as well as cyproterone acetate, azelaic acid and its derivatives and those compounds described in U.S. Patent Number 5,480,913, flutamide, and those compounds described in U.S. Patent Numbers 5,411,981, 5,565,467, and 4,910,226.

15 Antimicrobials include selenium sulfide, ketoconazole, triclocarbon, triclosan, zinc pyrithione, itraconazole, asiatic acid, hinokitiol, mipirocin and those described in EPA 0,680,745, clinacycin hydrochloride, benzoyl peroxide, benzyl peroxide and minocyclin.

Examples of suitable anti-inflammatories include glucocorticoids such as  
 20 hydrocortisone, mometasone furoate and prednisolone, nonsteroidal anti-inflammatories including cyclooxygenase or lipoxygenase inhibitors such as those described in U.S. Patent Number 5,756,092, and benzydamine, salicylic acid, and those compounds described in EPA 0,770,399, published May 2, 1997, WO 94/06434, published March 31, 1994, and FR 2,268,523, published November 21,  
 25 1975.

3,5,3'-Triiodothyronine is an example of a suitable thyroid hormone.

Examples of suitable non-selective prostaglandin agonists and antagonists include compounds such as those described in WO 98/33497, Johnstone, published August 6, 1998, WO 95/11003, Stjernschantz, published April 27, 1995, JP 97-  
 30 100091, Ueno and JP 96-134242, Nakamura.

Suitable retinoids include isotretinoin, acitretin, and tazarotene.

Other optional hair growth stimulants for component i) include

- benzalkonium chloride, benzethonium chloride, phenol, estradiol, chlorpheniramine maleate, chlorophyllin derivatives, cholesterol, salicylic acid, cysteine, methionine, red pepper tincture, benzyl nicotinate, D,L - menthol, peppermint oil, calcium
- 5 pantothenate, panthenol, castor oil, prednisolone, resorcinol, chemical activators of protein kinase C, glycosaminoglycan chain cellular uptake inhibitors, inhibitors of glycosidase activity, glycosaminoglycanase inhibitors, esters of pyroglutamic acid, hexosaccharic acids or acylated hexosaccharic acids, aryl-substituted ethylenes, N-
- 10 acylated amino acids, flavinoids, ascomycin derivatives and analogs, histamine antagonists such as diphenhydramine hydrochloride, triterpenes such as oleanolic acid and ursolic acid and those described in U.S. Patent Numbers 5,529,769, 5,468,888, 5,631,282, and 5,679,705, JP 10017431, WO 95/35103, JP 09067253, WO 92/09262, JP 62093215, and JP 08193094; saponins such as those described in
- 15 Bonte et al, published January 16, 1997, proteoglycanase or glycosaminoglycanase inhibitors such as those described in U.S. Patent Numbers 5,015,470, 5,300,284, and 5,185,325, estrogen agonists and antagonists, pseudoterins, cytokine and growth factor promoters, analogs or inhibitors such as interleukin1 inhibitors, interleukin-6 inhibitors, interleukin-10 promoters, and tumor necrosis factor inhibitors, vitamins
- 20 such as vitamin D analogs and parathyroid hormone antagonists, Vitamin B12 analogs and panthenol, interferon agonists and antagonists, hydroxyacids such as those described in U.S. Patent Number 5,550,158, benzophenones, and hydantoin anticonvulsants such as phenytoin, and combinations thereof.

- Other additional hair growth stimulants are described in JP 09-157,139 to
- 25 Tsuji et al., published June 17, 1997; EP 0277455 A1 to Mirabeau, published August 10, 1988; WO 97/05887 to Cabo Soler et al., published February 20, 1997; WO 92/16186 to Bonte et al., published March 13, 1992; JP 62-93215 to Okazaki et al., published April 28, 1987; U.S. Patent 4,987,150 to Kurono et al., issued January 22, 1991; JP 290811 to Ohba et al., published October 15, 1992; JP 05-286,835 to
- 30 Tanaka et al., published November 2, 1993, FR 2,723,313 to Greff, published August 2, 1994, U. S. Patent Number 5,015,470 to Gibson, issued May 14, 1991, U.S. Patent Number 5,559,092, issued September 24, 1996, U.S. Patent Number



5,536,751, issued July 16, 1996, U.S. Patent Number 5,714,515, issued February 3, 1998, EPA 0,319,991, published June 14, 1989, EPA 0,357,630, published October 6, 1988, EPA 0,573,253, published December 8, 1993, JP 61-260010, published November 18, 1986, U.S. Patent Number 5,772,990, issued June 30, 1998, U.S. Patent Number 5,053, 410, issued October 1, 1991, and U.S. Patent Number 4,761,401, issued August 2, 1988.

The most preferred activity enhancers are minoxidil and finasteride, most preferably minoxidil.

Component ii) is a penetration enhancer that can be added to all of the compositions for systemic administration. The amount of component ii), when present in the composition, is typically 1 to 5 %. Examples of penetration enhancers include 2-methyl propan-2-ol, propan-2-ol, ethyl-2-hydroxypropanoate, hexan-2,5-diol, polyoxyethylene(2) ethyl ether, di(2-hydroxypropyl) ether, pentan-2,4-diol, acetone, polyoxyethylene(2) methyl ether, 2-hydroxypropionic acid, 2-hydroxyoctanoic acid, propan-1-ol, 1,4-dioxane, tetrahydrofuran, butan-1,4-diol, propylene glycol dipelargonate, polyoxypropylene 15 stearyl ether, octyl alcohol, polyoxyethylene ester of oleyl alcohol, oleyl alcohol, lauryl alcohol, dioctyl adipate, dicapryl adipate, di-isopropyl adipate, di-isopropyl sebacate, dibutyl sebacate, diethyl sebacate, dimethyl sebacate, dioctyl sebacate, dibutyl suberate, dioctyl azelate, dibenzyl sebacate, dibutyl phthalate, dibutyl azelate, ethyl myristate, dimethyl azelate, butyl myristate, dibutyl succinate, didecyl phthalate, decyl oleate, ethyl caproate, ethyl salicylate, isopropyl palmitate, ethyl laurate, 2-ethyl-hexyl pelargonate, isopropyl isostearate, butyl laurate, benzyl benzoate, butyl benzoate, hexyl laurate, ethyl caprate, ethyl caprylate, butyl stearate, benzyl salicylate, 2-hydroxypropanoic acid, 2-hydroxyoctanoic acid, dimethyl sulphoxide, N,N-dimethyl acetamide, N,N-dimethyl formamide, 2-pyrrolidone, 1-methyl-2-pyrrolidone, 5-methyl-2-pyrrolidone, 1,5-dimethyl-2-pyrrolidone, 1-ethyl-2-pyrrolidone, phosphine oxides, sugar esters, tetrahydrofurfural alcohol, urea, diethyl-m-toluamide, 1-dodecylazacycloheptan-2-one, omega three fatty acids and fish oils, and combinations thereof.

In a preferred embodiment of the invention, the PGF's are topically administered. Topical compositions that can be applied locally to the skin may be in

any form including solutions, oils, creams, ointments, gels, lotions, shampoos, leave-on and rinse-out hair conditioners, milks, cleansers, moisturizers, sprays, skin patches, and the like. Topical compositions comprise: component A) the PGF described above and component B) a carrier. The carrier of the topical composition preferably aids penetration of the PGF's into the skin to reach the environment of the hair follicle. Component B) may further comprise one or more optional components. Topical compositions preferably further comprise C) one or more of the optional activity enhancers described above.

The exact amounts of each component in the topical composition depend on various factors. The amount of component A) depends on the  $IC_{50}$  of the PGF selected. "IC<sub>50</sub>" means inhibitory concentration 50<sup>th</sup> percentile. The amount of component A) added to the topical composition is:

$$IC_{50} \times 10^{-2} \geq \% \text{ of component A) } \geq IC_{50} \times 10^{-3},$$

where  $IC_{50}$  is expressed in nanomolar units. For example, if the  $IC_{50}$  of the PGF is 1 nM, the amount of component A) will be 0.001 to 0.01%. If the  $IC_{50}$  of the PGF is 10 nM, the amount of component A) will be 0.01 to 0.1%. If the  $IC_{50}$  of the PGF is 100 nM, the amount of component A) will be 0.1 to 1.0%. If the  $IC_{50}$  of the PGF is 1000 nM, the amount of component A) will be 1.0 to 10%, preferably 1.0 to 5%. If the amount of component A) is outside the ranges specified above (i.e., either higher or lower), efficacy of the treatment may be reduced.  $IC_{50}$  can be calculated according to the method in Reference Example 1, below. One skilled in the art can calculate  $IC_{50}$  without undue experimentation.

The topical composition preferably further comprises 1 to 20% component C), and a sufficient amount of component B) such that the amounts of components A), B), and C), combined equal 100%. The amount of B) the carrier employed in conjunction with the PGF is sufficient to provide a practical quantity of material for administration per unit dose of the compound. Techniques and compositions for making dosage forms useful in the methods of this invention are described in the following references: Modern Pharmaceutics, Chapters 9 and 10, Banker & Rhodes, eds. (1979); Lieberman et al., Pharmaceutical Dosage Forms: Tablets (1981); and Ansel, Introduction to Pharmaceutical Dosage Forms, 2<sup>nd</sup> Ed., (1976).

Component B) the carrier may comprise a single ingredient or a combination of two or more ingredients. In the topical compositions, component B) is a topical carrier. Preferred topical carriers comprise one or more ingredients selected from the group consisting of water, alcohols, aloe vera gel, allantoin, glycerin, vitamin A and E oils, mineral oil, propylene glycol, polypropylene glycol-2 myristyl propionate, dimethyl isosorbide, combinations thereof, and the like. More preferred carriers include propylene glycol, dimethyl isosorbide, and water.

The topical carrier may comprise one or more ingredients selected from the group consisting of q) emollients, r) propellants, s) solvents, t) humectants, u) thickeners, v) powders, and w) fragrances in addition to, or instead of, the preferred topical carrier ingredients listed above. One skilled in the art would be able to optimize carrier ingredients for the topical compositions without undue experimentation.

Ingredient q) is an emollient. The amount of ingredient q) in the topical composition is typically 5 to 95%. Suitable emollients include stearyl alcohol, glyceryl monoricinoleate, glyceryl monostearate, propane-1,2-diol, butane-1,3-diol, mink oil, cetyl alcohol, isopropyl isostearate, stearic acid, isobutyl palmitate, isocetyl stearate, oleyl alcohol, isopropyl laurate, hexyl laurate, decyl oleate, octadecan-2-ol, isocetyl alcohol, cetyl palmitate, di-n-butyl sebacate, isopropyl myristate, isopropyl palmitate, isopropyl stearate, butyl stearate, polyethylene glycol, triethylene glycol, lanolin, sesame oil, coconut oil, arachis oil, castor oil, acetylated lanolin alcohols, petrolatum, mineral oil, butyl myristate, isostearic acid, palmitic acid, isopropyl linoleate, lauryl lactate, myristyl lactate, decyl oleate, myristyl myristate, polydimethylsiloxane, and combinations thereof. Preferred emollients include stearyl alcohol and polydimethylsiloxane.

Ingredient r) is a propellant. The amount of ingredient r) in the topical composition is typically 5 to 95%. Suitable propellants include propane, butane, isobutane, dimethyl ether, carbon dioxide, nitrous oxide, and combinations thereof. Ingredient s) is a solvent. The amount of ingredient s) in the topical composition is typically 5 to 95 %. Suitable solvents include water, ethyl alcohol, methylene chloride, isopropanol, castor oil, ethylene glycol monoethyl ether, diethylene glycol monobutyl ether, diethylene glycol monoethyl ether, dimethylsulfoxide, dimethyl

formamide, tetrahydrofuran, and combinations thereof. Preferred solvents include ethyl alcohol.

Ingredient t) is a humectant. The amount of ingredient t) in the topical composition is typically 5 to 95 %. Suitable humectants include glycerin, sorbitol, sodium 2-pyrrolidone-5-carboxylate, soluble collagen, dibutyl phthalate, gelatin, and combinations thereof. Preferred humectants include glycerin.

Ingredient u) is a thickener. The amount of ingredient u) in the topical composition is typically 0 to 95%.

Ingredient v) is a powder. The amount of ingredient v) in the topical composition is typically 0 to 95 %. Suitable powders include chalk, talc, fullers earth, kaolin, starch, gums, colloidal silicon dioxide, sodium polyacrylate, tetra alkyl ammonium smectites, trialkyl aryl ammonium smectites, chemically modified magnesium aluminum silicate, organically modified montmorillonite clay, hydrated aluminum silicate, fumed silica, carboxyvinyl polymer, sodium carboxymethyl cellulose, ethylene glycol monostearate, and combinations thereof.

Ingredient w) is a fragrance. The amount of ingredient w) in the topical composition is typically 0.001 to 0.5%, preferably 0.001 to 0.1%.

Component C) the optional activity enhancer is as described above. Any of the i) hair growth stimulants and ii) penetration enhancers may be added to the topical compositions. Preferably, the topical composition comprises 0.01 to 15% of component i) the optional hair growth stimulant. More preferably, the composition comprises 0.1 to 10%, and most preferably 0.5 to 5% of component i). Preferably, the topical composition comprises 1 to 5% of component ii).

In an alternative embodiment of the invention, topical pharmaceutical compositions for ocular administration are prepared by conventional methods. Topical pharmaceutical compositions for ocular administration typically comprise A) a PGF, B) a carrier, such as purified water, and one or more ingredients selected from the group consisting of y) sugars such as dextrans, particularly dextran 70, z) cellulose or a derivative thereof, aa) a salt, bb) disodium EDTA (Edetate disodium), and cc) a pH adjusting additive.

Examples of z) cellulose derivatives suitable for use in the topical pharmaceutical composition for ocular administration include sodium

carboxymethyl cellulose, ethyl cellulose, methyl cellulose, and hydroxypropylmethylcellulose. Hydroxypropylmethylcellulose is preferred.

Examples of aa) salts suitable for use in the for use in the topical pharmaceutical composition for ocular administration include sodium chloride, potassium chloride, and combinations thereof.

Examples of cc) pH adjusting additives include HCl or NaOH in amounts sufficient to adjust the pH of the topical pharmaceutical composition for ocular administration to 7.2-7.5.

This invention further relates to a method for darkening hair, thickening hair, and reversing hair graying. The method comprises applying the topical composition for treating hair loss to hair, to skin in the locus of hair, or both. For example, the topical composition may be applied to hair growing on the scalp or eyelashes. The topical composition can be, for example, a cosmetic composition prepared as described above. An example of a composition that may be applied to eyelashes is a mascara. The prostaglandin may be added to mascara compositions known in the art, such as the mascara described in U.S. Patent No. 5,874,072, which is hereby incorporated by reference. The mascara comprises dd) a water-insoluble material, ee) a water-soluble, film-forming polymer, ff) a wax, o) a surfactant, gg) a pigment, and s) a solvent.

Ingredient dd) is a water-insoluble material selected from the group consisting of acrylate copolymers; styrene/acrylate/methacrylate copolymers; acrylic latex; styrene/acrylic ester copolymer latex; polyvinylacetate latex; vinyl acetate/ethylene copolymer latex; styrene/butadiene copolymer latex; polyurethane latex; butadiene/acrylonitrile copolymer latex; styrene/acrylate/acrylonitrile copolymer latex; and mixtures thereof, wherein the acrylate copolymers, and the styrene/acrylate/methacrylate copolymers additionally comprise ammonia, propylene glycol, a preservative and a surfactant.

Ingredient ee) is a water-soluble, film-forming polymer. Ingredient ee) is selected from the group consisting of vinyl alcohol/poly(alkyleneoxy)acrylate, vinyl alcohol/vinyl acetate/poly-(alkyleneoxy)acrylate, polyethylene oxide, polypropylene oxide, acrylates/octyl-acrylamide copolymers and mixtures thereof.

Ingredient ff) is a wax. "Wax" means a lower-melting organic mixture or compound of high molecular weight, solid at room temperature and generally similar in composition to fats and oils except that they contain no glycerides. Some are hydrocarbons, others are esters of fatty acids and alcohols. Waxes useful in this invention are selected from the group consisting of animal waxes, vegetable waxes, mineral waxes, various fractions of natural waxes, synthetic waxes, petroleum waxes, ethylenic polymers, hydrocarbon types such as Fischer-Tropsch waxes, silicone waxes, and mixtures thereof wherein the waxes have a melting point between 55 and 100°C.

Ingredient o) is surfactant, as described above. Ingredient o) in the mascara is preferably a surfactant having an HLB from 3 to 15. Suitable surfactants include those disclosed in the C.T.F.A. Cosmetic Ingredient Handbook, pp.587-592 (1992); Remington's Pharmaceutical Sciences, 15th Ed. pp. 335-337 (1975); and McCutcheon's Volume 1, Emulsifiers & Detergents, North American Edition, pp. 236-239 (1994).

Ingredient gg) is a pigment. Suitable pigments include inorganic pigments, organic lake pigments, pearlescent pigments, and mixtures thereof. Inorganic pigments useful in this invention include those selected from the group consisting of rutile or anatase titanium dioxide, coded in the Color Index under the reference CI 77,891; black, yellow, red and brown iron oxides, coded under references CI 77,499, 77,492 and, 77,491; manganese violet (CI 77,742); ultramarine blue (CI 77,007); chromium oxide (CI 77,288); chromium hydrate (CI 77,289); and ferric blue (CI 77,510) and mixtures thereof.

The organic pigments and lakes useful in this invention include those selected from the group consisting of D&C Red No. 19 (CI 45,170), D&C Red No. 9 (CI 15,585), D&C Red No. 21 (CI 45,380), D&C Orange No. 4 (CI 15,510), D&C Orange No. 5 (CI 45,370), D&C Red No. 27 (CI 45,410), D&C Red No. 13 (CI 15,630), D&C Red No. 7 (CI 15,850), D&C Red No. 6 (CI 15,850), D&C Yellow No. 5 (CI 19,140), D&C Red No. 36 (CI 12,085), D&C Orange No. 10 (CI 45,425), D&C Yellow No. 6 (CI 15,985), D&C Red No. 30 (CI 73,360), D&C Red No. 3 (CI

45,430) and the dye or lakes based on Cochineal Carmine (CI 75,570) and mixtures thereof.

The pearlescent pigments useful in this invention include those selected from the group consisting of the white pearlescent pigments such as mica coated with titanium oxide, bismuth oxychloride, colored pearlescent pigments such as titanium mica with iron oxides, titanium mica with ferric blue, chromium oxide and the like, titanium mica with an organic pigment of the above-mentioned type as well as those based on bismuth oxychloride and mixtures thereof.

Ingredient s) is a solvent described above, preferably water.

The amount of A) the PGF added to the mascara is as described above for topical compositions.

The PGF's may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines. A preferred formulation for topical delivery of the present compounds uses liposomes as described in Dowton et al., "Influence of Liposomal Composition on Topical Delivery of Encapsulated Cyclosporin A: I. An in vitro Study Using Hairless Mouse Skin", S.T.P. Pharma Sciences, Vol. 3, pp. 404 - 407 (1993); Wallach and Philippot, "New Type of Lipid Vesicle: Novasome<sup>®</sup>", Liposome Technology, Vol. 1, pp. 141 - 156 (1993); Wallach, U.S. Patent No. 4,911,928, assigned to Micro-Pak, Inc., issued March 27, 1990; and Weiner et al., U.S. Patent No. 5,834,014, assigned to The University of Michigan and Micro-Pak, Inc., issued November 10, 1998 (with respect to Weiner et al., with a compound as described herein administered in lieu of, or in addition to, minoxidil).

The PGF's may also be administered by iontophoresis. See, e.g., Internet site [www.unipr.it/arpa/dipfarm/erasmus/erasm14.html](http://www.unipr.it/arpa/dipfarm/erasmus/erasm14.html); Banga et al., "Hydrogel-based Iontotherapeutic Delivery Devices for Transdermal Delivery of Peptide/Protein Drugs", Pharm. Res., Vol. 10 (5), pp. 697-702 (1993); Ferry, "Theoretical Model of Iontophoresis Utilized in Transdermal Drug Delivery", Pharmaceutical Acta Helvetiae, Vol 70, pp. 279-287 (1995); Gangarosa et al., "Modern Iontophoresis for Local Drug Delivery", Int. J. Pharm, Vol. 123, pp. 159-171 (1995); Green et al.,

- “Iontophoretic Delivery of a Series of Tripeptides Across the Skin in vitro”, *Pharm. Res.*, Vol 8, pp. 1121-1127 (1991); Jadoul et al., “Quantification and Localization of Fentanyl and TRH Delivered by Iontophoresis in the Skin”, *Int. J. Pharm.*, Vol. 120, pp. 221-8 (1995); O'Brien et al., “An Updated Review of its Antiviral Activity, Pharmacokinetic Properties and Therapeutic Efficacy”, *Drugs*, Vol. 37, pp. 233-309 (1989); Parry et al., “Acyclovir Bioavailability in Human Skin”, *J. Invest. Dermatol.*, Vol. 98 (6), pp. 856-63 (1992); Santi et al., “Drug Reservoir Composition and Transport of Salmon Calcitonin in Transdermal Iontophoresis”, *Pharm. Res.*, Vol 14 (1), pp. 63-66 (1997); Santi et al., “Reverse Iontophoresis - Parameters Determining Electroosmotic Flow: I. pH and Ionic Strength”, *J. Control. Release*, Vol. 38, pp. 159-165 (1996); Santi et al., “Reverse Iontophoresis - Parameters Determining Electroosmotic Flow: II. Electrode Chamber Formulation”, *J. Control. Release*, Vol. 42, pp. 29-36 (1996); Rao et al., “Reverse Iontophoresis: Noninvasive Glucose Monitoring in vivo in Humans”, *Pharm. Res.*, Vol. 12 (12), pp. 1869-1873 (1995); Thysman et al., “Human Calcitonin Delivery in Rats by Iontophoresis”, *J. Pharm. Pharmacol.*, Vol. 46, pp. 725-730 (1994); and Volpato et al., “Iontophoresis Enhances the Transport of Acyclovir through Nude Mouse Skin by Electrorepulsion and Electroosmosis”, *Pharm. Res.*, Vol. 12 (11), pp. 1623-1627 (1995).

The PGF's may be included in kits comprising a PGF, a systemic or topical composition described above, or both; and information, instructions, or both that use of the kit will provide treatment for hair loss in mammals (particularly humans). The information and instructions may be in the form of words, pictures, or both, and the like. In addition or in the alternative, the kit may comprise a PGF, a composition, or both; and information, instructions, or both, regarding methods of application of the PGF or composition, preferably with the benefit of treating hair loss in mammals.

In all of the foregoing compositions, and for all routes of administration, the PGF's can be used alone or in combinations of two or more PGF's. The compositions may further comprise additional drugs or excipients as appropriate for the indication.

### Methods of the Invention



This invention further relates to a method for treating hair loss in mammals. The method comprises administering to a mammal (preferably a human) suffering from hair loss, a PGF described above. For example, a mammal diagnosed with alopecia including male pattern baldness and female pattern baldness can be treated  
 5 by the methods of this invention. Preferably, a systemic or topical composition comprising A) the PGF and B) a carrier is administered to the mammal. More preferably, the composition is a topical composition comprising A) the PGF, B) the carrier, and C) an optional activity enhancer.

The dosage of the PGF administered depends on the method of  
 10 administration. For systemic administration, (e.g., oral, rectal, nasal, sublingual, buccal, or parenteral), typically, 0.5 mg to 300 mg, preferably 0.5 mg to 100 mg, more preferably 0.1 mg to 10 mg, of a PGF described above is administered per day. These dosage ranges are merely exemplary, and daily administration can be adjusted depending on various factors. The specific dosage of the PGF to be  
 15 administered, as well as the duration of treatment, and whether the treatment is topical or systemic are interdependent. The dosage and treatment regimen will also depend upon such factors as the specific PGF used, the treatment indication, the efficacy of the compound, the personal attributes of the subject (such as, for example, weight, age, sex, and medical condition of the subject), compliance  
 20 with the treatment regimen, and the presence and severity of any side effects of the treatment.

For topical administration (e.g., local application on the skin, ocular, liposome delivery systems, or iontophoresis), the topical composition is typically administered once per day. The topical compositions are administered daily for a  
 25 relatively short amount of time (i.e., on the order of weeks). Generally, 6 to 12 weeks is sufficient. The topical compositions are preferably leave-on compositions. In general, the topical composition should not be removed for at least several hours after administration.

In addition to the benefits in treating hair loss, the inventors have surprisingly  
 30 found that the PGF's in the compositions and methods of this invention also darken and thicken hair and may reverse hair graying.

## EXAMPLES

These examples are intended to illustrate the invention to those skilled in the art and should not be interpreted as limiting the scope of the invention set forth in the claims.

5

### Reference Example 1 - Radioligand Binding Assay

IC<sub>50</sub> of a PGF can be determined relative to PGF<sub>2α</sub> using the Radioligand Binding Assay. As a control, the IC<sub>50</sub> for PGF<sub>2α</sub> itself should be no lower than 1.0 nM and no higher than 5.0 nM.

10

In this assay, COS-7 cells are transiently transfected with the hFP recombinant plasmid using LipofectAMINE Reagent. Forty-eight hours later, the transfected cells are washed with Hank's Balanced Salt Solution (HBSS, without CaCl<sub>2</sub>, MgCl<sub>2</sub>, MgSO<sub>4</sub>, or phenol red). The cells are detached with versene, and HBSS is added. The mixture is centrifuged at 200g for 10

15

minutes, at 4°C to pellet the cells. The pellet is resuspended in Phosphate-Buffered Saline-EDTA buffer (PBS; 1 mM EDTA; pH 7.4; 4°C). The cells are disrupted by nitrogen cavitation (Parr model 4639), at 800 psi, for 15 minutes at 4°C. The mixture is centrifuged at 1000g for 10 minutes at 4°C.

20

The supernatant is centrifuged at 100,000g for 60 minutes at 4°C. The pellet is resuspended to 1 mg protein/mL TME buffer (50 mM Tris; 10 mM MgCl<sub>2</sub>; 1 mM EDTA; pH 6.0; 4°C) based on protein levels measured using the Pierce BCA Protein Assay kit. The homogenate is mixed for 10 seconds using a Kinematica POLYTRON® (available from KINEMATICA AG, Luzernerstrasse 147A CH-6014 Littau, Switzerland). The membrane

25

preparations are then stored at -80°C, until thawed for assay use.

The receptor competition binding assays are developed in a 96 well format. Each well contains 100 g of hFP membrane, 5 nM (3 H) PGF<sub>2</sub>, and the various competing compounds in a total volume of 200 L. The plates are incubated at 23°C for 1 hour. The incubation is terminated by rapid filtration using the Packard

30

Filtermate 196 harvester through Packard UNIFILTER® GF/B filters (available from Packard Instrument Co., Inc. of Downers Grove Illinois) pre-wetted with TME

buffer. The filter is washed four times with TME buffer. Packard Microscint 20, a high efficiency liquid scintillation cocktail, is added to the filter plate wells and the plates remain at room temperature for three hours prior to counting. The plates are read on a Packard TOPCOUNT® Microplate Scintillation Counter (also available  
 5 from Packard Instrument Co., Inc.)

#### Reference Example 2 - Telogen Conversion Assay

PGF's are tested for their potential to grow hair using the Telogen Conversion Assay. The Telogen Conversion Assay measures the potential of a PGF  
 10 to convert mice in the resting stage of the hair growth cycle ("telogen"), to the growth stage of the hair growth cycle ("anagen").

Without intending to be limited by theory, there are three principal phases of the hair growth cycle: anagen, catagen, and telogen. It is believed that there is a longer telogen period in C3H mice (Harlan Sprague Dawley, Inc., Indianapolis, IN)  
 15 from approximately 40 days of age until about 75 days of age, when hair growth is synchronized. It is believed that after 75 days of age, hair growth is no longer synchronized. Wherein about 40 day-old mice with dark fur (brown or black) are used in hair growth experiments, melanogenesis occurs along with hair (fur) growth wherein the topical application of hair growth inducers are evaluated. The Telogen  
 20 Conversion Assay herein is used to screen PGF's for potential hair growth by measuring melanogenesis.

Three groups of 44 day-old C3H mice are used: a vehicle control group, a positive control group, and a test PGF group, wherein the test PGF group is administered a PGF used in the method of this invention. The length of the assay is  
 25 24 days with 15 treatment days (wherein the treatment days occur Mondays through Fridays). Day 1 is the first day of treatment. A typical study design is shown in Table 1 below. Typical dosage concentrations are set forth in Table 1, however the skilled artisan will readily understand that such concentrations may be modified.

30 Table 1 - Assay Parameters

Group #	Animal #	Compound	Concentration	Application	Length
---------	----------	----------	---------------	-------------	--------

				volume	of Study
1	1 - 10	Test Compound	0.01% in vehicle**	400 $\mu$ L topical	26 days
2	11 - 20	Positive Control (T3)*	0.01% in vehicle**	400 $\mu$ L topical	26 days
3	21 - 30	Vehicle**	N/A	400 $\mu$ L topical	26 days

\* T3 is 3,5,3'-triiodothyronine.

\*\*The vehicle is 60% ethanol, 20% propylene glycol, and 20% dimethyl isosorbide (commercially available from Sigma Chemical Co., St. Louis, MO).

- 5 The mice are treated topically Monday through Friday on their lower back (base of tail to the lower rib). A pipettor and tip are used to deliver 400  $\mu$ L to each mouse's back. The 400  $\mu$ L application is applied slowly while moving hair on the mouse to allow the application to reach the skin.


- 10 While each treatment is being applied to the mouse topically, a visual grade of from 0 to 4 will be given to the skin color in the application area of each animal. As a mouse converts from telogen to anagen, its skin color will become more bluish-black. As indicated in Table 2, the grades 0 to 4 represent the following visual observations as the skin progresses from white to bluish-black.

15 Table 2 - Evaluation Criteria

<u>Visual Observation</u>	<u>Grade</u>
Whitish Skin Color	0
Skin is light gray (indication of initiation of anagen)	1
Appearance of Blue Spots	2
Blue Spots are aggregating to form one large blue area	3
Skin is dark blue (almost black) with color covering majority of treatment area (indication of mouse in full anagen)	4

Example 1


A PGF having the structure:



was tested according to the method of Reference Example 1. The average grade was calculated by averaging the grades of 7 mice after 23 days, 25 days, and 26 days. The results are in Table 3. The PGF grew hair.

### Example 2

A PGF having the structure:



was tested according to the method of Reference Example 1. The average grade was calculated by averaging the grades of 7 mice after 23 days. The results are in Table 3. The PGF grew hair.

Table 3 - Average Grades

Example	23 Days	25 Days	26 Days
1	0.4	0.1	0.7
2	0.1	not measured	not measured

### Example 3

Compositions for topical administration are made, comprising:

Component	3-1	3-2	3-3
PGF (wt %)	0.07	0.1	2.4

IC <sub>50</sub> the PGF (nM)	7	10	24
Ethanol (wt %)	59.958	59.9	58.6
Propylene Glycol (wt %)	19.986	20.0	19.5
Dimethyl Isosorbide (wt %)	19.986	20.0	19.5

The PGFs in the compositions are as follows:

Sample	PGF
3-1	
3-2	
3-3	

A human male subject suffering from male pattern baldness is treated by a method of  
5 this invention. Specifically, for 6 weeks, one of the above compositions is daily  
administered topically to the subject to induce hair growth.

Example 4

A composition for topical administration is made according to the method of Dowton et al., "Influence of Liposomal Composition on Topical Delivery of Encapsulated Cyclosporin A: I. An in vitro Study Using Hairless Mouse Skin", S.T.P. Pharma Sciences, Vol. 3, pp. 404 - 407 (1993), using a PGF in lieu of cyclosporin A and using the NOVASOME® 1 (available from Micro-Pak, Inc. of Wilmington, Delaware) for the non-ionic liposomal formulation.

A human male subject suffering from male pattern baldness is treated each day with the above composition. Specifically, for 6 weeks, the above composition is administered topically to the subject.

Example 5

Shampoos are made, comprising:

Component	Ex. 5-1	Ex. 5-2	Ex. 5-3	Ex. 5-4
Ammonium Lauryl Sulfate	11.5 %	11.5 %	9.5 %	7.5 %
Ammonium Laureth Sulfate	4 %	3 %	2 %	2 %
Cocamide MEA	2 %	2 %	2 %	2 %
Ethylene Glycol Distearate	2 %	2 %	2 %	2 %
Cetyl Alcohol	2 %	2 %	2 %	2 %
Stearyl Alcohol	1.2 %	1.2 %	1.2 %	1.2 %
Glycerin	1 %	1 %	1 %	1 %
Polyquaternium 10	0.5 %	0.25 %	-	-
Polyquaternium 24	-	-	0.5 %	0.25 %
Sodium Chloride	0.1 %	0.1 %	0.1 %	0.1 %
Sucrose Polyesters of Cottonate Fatty Acid	3 %	3 %	-	-
Sucrose Polyesters of Behenate Fatty Acid	2 %	3 %	-	-
Polydimethyl Siloxane	-	-	3 %	2 %
Cocaminopropyl Betaine	-	1 %	3 %	3 %
Lauryl Dimethyl Amine Oxide	1.5 %	1.5 %	1.5 %	1.5 %
Decyl Polyglucose	-	-	1 %	1 %
DMDM Hydantoin	0.15 %	0.15 %	0.15 %	0.15 %
PGF having IC <sub>50</sub> of 7 nM	-	0.007 %	0.007 %	-
PGF having IC <sub>50</sub> of 24 nM	0.024 %	-	-	0.024 %
Minoxidil			3 %	2 %
Phenoxyethanol	0.5 %	0.5 %	0.5 %	0.5 %
Fragrance	0.5 %	0.5 %	0.5 %	0.5 %
Water	q.s.	q.s.	q.s.	q.s.

- A human subject suffering from male pattern baldness is treated by a method
- 5 of this invention. Specifically, for 12 weeks, a shampoo described above is used daily by the subject.



**Example 6**

A mascara composition is prepared. The composition comprises:

Component	% W/W
WATER, DEIONIZED, USP	q.s.
BLACK 1080 MICRONIZED TYPE	10.000
GLYCERYL MONOSTEARATE (2400 TYPE)	8.500
C18-36 ACID TRIGLYCERIDE	5.500
STEARIC ACID, TRIPLE PRESSED, LIQUID	4.000
ETHYL ALCOHOL SD 40-B, 190 PROOF/SERIAL #:	4.000
BEEWAX WHITE, FLAKES	3.250
SHELLAC, NF	3.000
LECITHIN, GRANULAR (TYPE 6450)	2.500
TRIETHANOLAMINE 99% - TANK	2.470
PARAFFIN WAX	2.250
PARAFFIN WAX 118/125	2.250
CARNAUBA WAX, NF	2.000
POTASSIUM CETYL PHOSPHATE	1.000
PHENOXYETHANOL	0.800
OLEIC ACID NF	0.750
DL-PANTHENOL	0.350
PVP/VA COPOLYMER	0.250
METHYL PARABEN, NF	0.200
DIAZOLIDINYL UREA	0.200
SIMETHICONE	0.200
ETHYL PARABEN NF	0.150
PENTAERYTHRITYL HYDROGENATED ROSINATE	0.150
PROPYL PARABEN, NF	0.100
TRISODIUM EDTA	0.100
PGF having IC <sub>50</sub> of 7 nM	0.007

The PGF is the same as that in Example 3-1.

- 5 A human female subject applies the composition each day. Specifically, for 6 weeks, the above composition is administered topically to the subject to darken and thicken eyelashes.

**Example 7**

- 10 Pharmaceutical compositions in the form of tablets are prepared by conventional methods, such as mixing and direct compaction, formulated as follows:

Ingredient	Quantity (mg per tablet)
PGF	0.5
Microcrystalline Cellulose	100
15 Sodium Starch Glycollate	30
Magnesium Stearate	3

The prostaglandin is the same as that in Example 3-3.

The above composition is administered orally to a subject once daily for 6 to 12 weeks to promote hair growth.

5

#### Example 8

Pharmaceutical compositions in liquid form are prepared by conventional methods, formulated as follows:

	<u>Ingredient</u>	<u>Quantity</u>
10	PGF	0.1 mg
	Phosphate buffered physiological saline	10 ml
	Methyl Paraben	0.05ml

The prostaglandin is the same as that in Example 3-3.

1.0 ml of the above composition is administered subcutaneously at the site of hair loss once daily for 6 to 12 weeks to promote hair growth.

15

#### Example 9

A topical pharmaceutical composition is prepared by conventional methods and formulated as follows:

	<u>Ingredient</u>	<u>Amount (wt %)</u>
20	PGF	0.004
	Dextran 70	0.1
	Hydroxypropyl methylcellulose	0.3
	Sodium Chloride	0.77
25	Potassium chloride	0.12
	Disodium EDTA (Edetate disodium)	0.05
	Benzalkonium chloride	0.01
	HCL and/or NaOH	pH 7.2-7.5
	Purified water	q.s. to 100%

30

The prostaglandin is the same as that in Example 3-3.


The above composition is administered ocularly to a subject once per day for 6 to 12 weeks to promote eyelash growth.

#### Effects of the Invention

- 5           The compositions and methods herein provide a cosmetic benefit with respect to hair growth and appearance in subjects desiring such treatment.
- What is claimed is:

1. A topical composition for treating hair loss comprising:

A) an active ingredient selected from the group consisting of a prostaglandin F analog having the structure



and pharmaceutically acceptable salts and hydrates of the prostaglandin F analog; biohydrolyzable amides, esters, and imides of the prostaglandin F analog; optical isomers, diastereomers, and enantiomers of the prostaglandin F analog; and combinations thereof;

wherein  $R^1$  is selected from the group consisting of  $\text{CO}_2\text{H}$ ,  $\text{C}(\text{O})\text{NHOH}$ ,  $\text{CO}_2\text{R}^5$ ,  $\text{CH}_2\text{OH}$ ,  $\text{S}(\text{O})_2\text{R}^5$ ,  $\text{C}(\text{O})\text{NHR}^5$ ,  $\text{C}(\text{O})\text{NHS}(\text{O})_2\text{R}^5$ , and tetrazole;

$R^2$  is selected from the group consisting of a hydrogen atom and a lower monovalent hydrocarbon group;

$R^3$  and  $R^4$  are each independently selected from the group consisting of H,  $\text{CH}_3$ ,  $\text{C}_2\text{H}_5$ ,  $\text{OR}^{10}$ ,  $\text{SR}^{10}$ , and OH; with the proviso that both  $R^3$  and  $R^4$  are not OH;

$R^5$  is selected from the group consisting of monovalent hydrocarbon groups, substituted monovalent hydrocarbon groups, aromatic groups, substituted aromatic groups, carbocyclic groups, substituted carbocyclic groups, heterogeneous groups, substituted heterogeneous groups, heterocyclic groups, substituted heterocyclic groups, heteroaromatic groups, and substituted heteroaromatic groups;

X is selected from the group consisting of  $\text{NR}^6\text{R}^7$ ,  $\text{OR}^8$ ,  $\text{SR}^9$ ,  $\text{S}(\text{O})\text{R}^9$ , and  $\text{S}(\text{O})_2\text{R}^9$ ;

$R^6$ ,  $R^7$ , and  $R^8$  are each independently selected from the group consisting of hydrogen atoms, acyl groups, monovalent hydrocarbon groups, substituted monovalent hydrocarbon groups, heterogeneous groups, substituted heterogeneous groups, carbocyclic groups, substituted carbocyclic groups, aromatic groups,